



UDC 637.146.3.05/.07-049.34

DOI: 10.31548/animal2021.03.009

The impact of storage on the sensory, viscosity and chemical characteristics of yogurt fortified with oat β -glucan

Xiaoqing Qu^{1,2}

¹Sumy National Agrarian University
40021, Herasima Kondratieva St. 160, Sumy, Ukraine

²Henan Institute of Science and Technology
453003, HuaLan Avenue, Xinxiang, China

Abstract. Oat β -glucan has the potential of a functional additive to yogurt, however, texture analysis and chemical characteristics during storage of yogurt with the addition of this additive require more systematic research, which determines the relevance of the study. The aim of the research work was to analyze the feasibility of enriching yogurt with 0.3% oat β -glucan. A number of methods for obtaining new knowledge in the direction of developing healthy food products were used. It was found that yogurt with the addition of 0.3% oat β -glucan had water-holding capacity. All pH values ranged from 4.18 to 4.28, which are within the normal ranges for set-type yogurts. A higher viscosity of the experimental sample of yogurt compared to the control was established, and the highest values were 58560 ± 2120 cp at 21 days for yogurt with the addition of 0.3% oat β -glucan. The viability of probiotic bacteria in yogurts was examined. During the whole storage period, the content of probiotics decreased, but yogurt with the addition of 0.3% oat β -glucan contained significantly more live probiotic bacteria compared to the control. The viability of probiotic bacteria of 0.3% oat β -glucan yogurt at 14 d ($3.18 \pm 0.2 \times 10^7$ CFU/mL) was only slightly lower than that of control yogurt at 1 d ($3.45 \pm 0.3 \times 10^7$ CFU/mL). This fully demonstrates that the addition of 0.3% oat β -glucan has a protective effect on probiotics in yogurt, which will be beneficial for human health. It was found that the addition of 0.3% oat β -glucan led to a decrease in adhesiveness, but an increase in hardness and stickiness during storage. The results of the sensory evaluation showed the highest acceptability values of 86.49 for 21 days of storage of yogurt with the addition of 0.3% oat β -glucan. The practical value of the research work lies in the developed yogurt containing 0.3% oat β -glucan being possibly to use as an innovative healthy dairy product

Keywords: yogurt, oat β -glucan, sensory, viscosity, chemical characteristics

Suggested Citation:

Qu, X. (2021). The impact of storage on the sensory, viscosity and chemical characteristics of yogurt fortified with oat β -glucan. *Animal Science and Food Technology*, 12(3), 96-104.

*Corresponding author

Introduction

The demand for healthy foods has caused the development of products containing functional components. In the dairy industry, prebiotic substances or probiotic bacteria have sparked interest due to scientific evidence related to their positive health benefits [1].

Dietary fibers, usually polysaccharides, can be insoluble or soluble in water. Soluble fibers are usually found in fruits, vegetables, oat bran, and barley, etc. Oat β -glucan (OG), a soluble dietary fiber from oat, has received tremendous attention during the previous decades. It is a non-starchy polysaccharide that comprises mainly glucose molecules linked by (1-3), (1-4) mixed linkage with glycosidic bonds [2]. More importantly, it has diverse and great effects on the maintenance and improvement of human health. For example, consumption of OG can help in reducing the occurrence of some chronic diseases such as high blood pressure, insulin responses, obesity, and colon cancer [3]. In addition, it can modify the rheological and mouthfeel attributes, and physical stability of beverages, and be used as a prebiotic in yogurt formulations [4]. Nowadays, consumption of OG is encouraged and it is added to various food products as a functional ingredient.

Analysis of Recent Researches and Publications

Yogurt, a traditional fermented dairy product containing probiotics and prebiotics, has gained widespread consumer acceptance for being considered as a healthy product [5, 6]. It has shown a substantial consumption increase in recent years due to its nutritional and health interest. During the fermentation action, some toxic or antinutritional factors, such as lactose and galactose, can be removed from fermented milk, thus preventing lactose intolerance and accumulation of galactose [7]. In this sense,

yogurt can be used as an excellent matrix for developing innovative health-promoting products and functional foods.

The addition of soluble fibers can change the structural-mechanical properties [8] and functional characteristics of fermented dairy products. For example, used hydrolyzed guar gum can be used to enhance the functional and sensory properties of yogurt [9]. In recent years, there is growing interest to launch health-promoting foodstuffs with OG.

Because OG is a kind of dietary fiber, it can be used as a prebiotic of probiotics, and it has a positive influence on the physicochemical properties and vitality of probiotics in yogurt. This feature is conducive to OG as a functional component of yogurt [10, 11].

From the analysis data of literature sources, it can be said that OG has the potential as a functional additive for yogurt; however, the texture analysis and chemical characteristics of the storage period after adding OG to yogurt still need to be more systematically studied.

The addition of 0.3% OG decreased the fermentation time of set-type yogurt, making the highest score of sensory evaluation [12]. As a functional food ingredient, 0.3% OG is important in enhancing the nutraceutical quality and physical characteristics of yogurt. Moreover, 0.3% OG shows its effect on starter culture growth. However, the impact of 0.3% OG on the quality of yogurt, such as sensory evaluation and physical characteristics, during storage, has not been evaluated. Therefore, in this work, the quality characteristics of set-type yogurt, i.e., control yogurt and 0.3% OG yogurt, were studied. The water-holding capacity (WHC), acidity value, pH, viscosity, textural parameters, viability of probiotic bacteria, and sensory properties, during 21 d of storage, were evaluated and compared.

The aim and objectives of the study. The aim of the study is to analyze the effect of the

feasibility of set-type yogurt fortification with 0.3% OG.

To achieve the aim of the study, it was necessary to solve the following problems:

- to analyze the effects of yogurt with or without 0.3% OG on sensory during storage;
- to examine the impact of yogurt with or without 0.3% OG on viscosity and chemical properties during storage;
- to investigate the effect on probiotics in yogurt with or without 0.3% OG during storage.

Materials and Methods

Pure milk was purchased from Yili Industrial Group Co. Ltd (Neimenggu, China). Oat β -glucan (95% purity) was provided by Zhongkang Food Co., (Guangzhou, China). Starters: *Streptococcus thermophilus* and *Lactobacillus bulgaricus* (*Lactobacillus dechellii* Bulgarian subspecies) (viable bacteria count was about 1×10^9 CFU/g) were provided by Danisco (China) Co., Ltd, (Shanghai, China). MRS medium was purchased from Shanghai Mingrui Biotechnology Co., Ltd, (Shanghai, China). Agar was purchased from Xi'an Huibang Bioengineering Co., Ltd, (Xi'an, China). All other chemicals were used of analytical grade.

The preparation of yogurt samples refers to the previous experimental methods [12].

Take a certain quality (M_1) of yogurt in a centrifuge tube; centrifuge at 4 000 r/min for 25 min at 4°C, discard the supernatant and weigh the remaining mass (M_2). It is not easy to separate whey from yogurt with good water-holding capacity during storage [13].

The calculation formula is as follows:

$$WHC (\%) = M_2 \div M_1 \times 100\% \quad (1)$$

Determination of acidity by an acidimeter (PB-10, Seidolis instruments, Germany): set-type yogurt samples were taken every 1 h during the fermentation process and the yogurt samples were determined after being stored

24 h. Then, 10 g yogurt samples were put into a 250 mL triangle bottle and 20 mL distilled water was added to dilute and mixed, 0.5% phenolphthalein was added as an indicator and titrated with 0.1 mol/L NaOH standard solution until it was slightly red. Not fading within 30 s marked the end of determination. Consuming 0.1 M of NaOH is equal to 1 °T.

All pH values were monitored after adding starter bacterial cultures using a digital pH meter (Thermol Scientific Inc., USA).

The viscosity of yogurt was measured at 4°C with a spindle (No. 3) rotation of 1500 rpm using the digital display rotating viscometer (NDJ-8S, Shanghai Yueping Scientific Instrument Co., Ltd., Shanghai, China). The readings were recorded at the 20th s of the measurement. The measurements were made in triplicate.

The experiments were performed by a texture analyzer (TA-Xt.plus Texture Analyzer; Stable Micro Systems, Godalming, UK). Refer to Bedani *et al.*, Ciron *et al.* (2010) and other methods, slightly improved. The yogurt samples were placed in 25 mL beakers and the Texture Profile Analysis (TPA) test mode and P/0.5 probe were used to determine the gel structure of the fermented milk. The parameters are as follows: the speed is divided into 2.0 mm/s before measurement, 1.0 mm/s during measurement, and 2.0 mm/s after measurement. Test samples shall be made in triplicate.

Microbial Analyses. The 1 g of yogurt with 9 mL of 0.9% (w/v) NaCl was mixed and diluted to a concentration of 10^4 , 10^5 , and 10^6 , and then 1 mL of each dilution was inoculated on plates containing the MRS agar. Bacteria were counted by the pour plate technique. The plates in duplicates were incubated anaerobically at 37°C for 72 h, and then colonies were counted [4, 14, 15].

According to the relevant sensory evaluation methods of the national standard, 10 volunteers (5 men and 5 women) with sensory

evaluation experience of fermented milk were selected to conduct the sensory evaluation of fermented milk added with OG from the aspects of color, structural state, texture, and flavor. The full score was 100 and the sensory score was the average score of 10 people.

Unless otherwise stated, all tests were performed in triplicate, and data were averaged. Standard deviation was also calculated. SPSS version 17.0 (SPSS Inc., Chicago, IL, USA) was used for all statistical evaluations, and OriginPro 8.6.0 (Originlab, Northampton, Massachusetts, USA) was used for the construction of the graphs. The sensory evaluation and yogurt WHC with different storage times of OG were analyzed by independent sample t-test of SPSS 17.0 software. The significance (P values) of the

microbiological parameters were calculated by the general linear model. One-way ANOVA was used to analyze the titration acidity of yogurt for different storage time, LSD and Dunnett's T3 test were applied for multiple comparisons and differences were considered to be statistically significant at $P < 0.05$.

Results of the Research and Discussion

For water-holding capacity values, a significant decrease was observed in the control yogurt, while a slight decrease was observed in 0.3% OG yogurt (Fig. 1 (a)). This can be attributed to the hydration properties of OG, which have an impact on the shelf life of food products by preventing texture loss and avoiding syneresis problems.




Figure 1. The WHC (a), acidity value (b), pH (c), and viscosity (d) of yogurt

As can be seen from Figure 1(b), the acidity of both control yogurt and 0.3% OG yogurt significantly increased to 93.8 ± 1.2 during 7 d of storage. And then (7-21 d of storage), no significant ($P \leq 0.05$) changes were observed in acidity values. An opposite trend was observed for pH values and reached the minimum (about 4.21 in

control yogurt and about 4.18 in 0.3% OG yogurt) (Fig. 1(c)). The decrease in pH values during the storage period might be mainly due to the utilization of OG by viable probiotic bacteria and the production of lactic acid. In addition, small amounts of CO_2 and formic acid from lactose may also lead to a decrease in pH [16, 17].

All pH values ranged from 4.18 to 4.28, which are within the normal ranges for set-type yogurts.

Figure 1(d) shows the changes in viscosity values over 21 d of storage. Significant differences were noted between control yogurt and 0.3% OG yogurt. In general, the viscosity values increased throughout storage in concentrated and non-fat plain yogurt throughout storage [18, 19]. The increasing viscosity during storage could be due to the protein rearrangement and protein-protein contact. The 0.3% OG yogurt had a higher viscosity than the control yogurt, and the highest values were 58560 ± 2120 cp at 21 d for 0.3% OG yogurt. This change could be attributed to the fact that OG has the ability to entrap water within the product. On the other hand, the addition of 0.3% OG could improve the protein rearrangement and protein-protein contact.




Figure 2. The viability of probiotic bacteria in yogurts during the storage

Live and active probiotic bacteria are considered to be beneficial for human health [20, 21]. OG is considered to be prebiotic and may contribute to the activity of probiotic bacteria. So, the viability of probiotic bacteria in yogurts was checked. As shown in Figure 2, during the whole storage period, the content of probiotics decreased, which was only $0.63 \pm 0.05 \times 10^7$ CFU/mL at 21 d. However, 0.3% OG yogurt contained significantly more live probiotic bacteria compared to the control one ($P < 0.05$) throughout the whole cold storage period. The viability of probiotic bacteria of 0.3% OG yogurt at 14 d ($3.18 \pm 0.2 \times 10^7$ CFU/mL) was only slightly lower than that of the control yogurt at 1 d ($3.45 \pm 0.3 \times 10^7$ CFU/mL). Similar results have also been reported [15]. This fully demonstrates that the addition of OG has a protective effect on probiotics in yogurt. The survival of yogurt microbiota in 0.3% OG yogurt will be beneficial for human health.

The textural parameters of yogurt are important for yogurt products, as they can simulate their breakdown occurring in the mouth. Results of texture profile analysis (TPA) of yogurts, including hardness (N), consistency, cohesiveness, and springiness (%) are summarized in Table 1. For both control yogurt and 0.3% OG yogurt, the hardness of yogurts was improved, while the adhesiveness and gumminess values declined during the storage. The textural characteristics of yogurts are affected by the addition of 0.3% OG.

Table 1. Results of TPA test in the control yogurt and 0.3% OG yogurt

Test	Days	Yogurt samples	
		control yogurt	0.3% OG yogurt
Hardness	1	21.14	22.85
	7	24.77	28.36
	14	28.76	35.99
	21	40.99	50.45
Adhesiveness	1	-8.23	-9.15
	7	-9.18	-10.21

Table 1. Continued

Test	Days	Yogurt samples	
		control yogurt	0.3% OG yogurt
Gumminess	14	-9.25	-11.37
	21	-10.39	-11.85
	1	9.26	13.78
	7	12.79	13.81
Cohesiveness	14	14.71	19.82
	21	21.54	13.78
	1	0.43	0.43
	7	0.41	0.44
Chewiness	14	0.51	0.57
	21	0.53	0.59
	1	8.75	9.46
	7	11.39	12.87
Springiness	14	14.06	13.77
	21	20.72	18.99
	1	0.94	0.92
	7	0.95	0.93
	14	0.95	0.95
	21	0.96	0.97

The evaluation index of yogurt is mainly the hardness. The hardness of 0.3% OG yogurt was higher than that of control yogurt, especially after 21 days of storage (50.45 in 0.3% OG yogurt and 40.99 in the control yogurt). The higher hardness values could be attributed to the ability of the OG to entrap water, and maintain structure within the product [19, 22].

In contrast, the adhesiveness of 0.3% OG yogurt was lower than that of the control yogurt, especially after 21 d of storage (-11.85 in 0.3% OG yogurt and -10.39 in the control yogurt). The decrease in adhesiveness may be due to the fact that OG can reduce adherence of the yogurt with teeth during chewing.

The 0.3% OG yogurt had a slight increase in gumminess as compared to the control yogurt. And, during storage, an increase in gumminess was observed, which might be due to the increased hardness of yogurts. Similar behavior was found for cohesiveness, which can be attributed to the fact that OG has the ability to form new gel structures with casein, effectively

intercepting and entrapping water within the yogurt. No significant differences were obtained for springiness values between the control yogurt and 0.3% OG yogurt.

In short, results of textural profile tests showed appropriate hardness, less gumminess, less adhesiveness, and less destruction in 0.3% OG yogurt, giving a firm and creamy texture that was near to creamy mouthfeel of full-fat yogurt, improving the mouthfeel characteristics thereby enhanced sensory appeal for the product in which incorporated. The sensory properties of the yogurts are shown in Table 2. All yogurts had average sensory scores of above 80, indicating a preference both for the control yogurt and 0.3% OG yogurt throughout storage.

The sensory results indicated that there was a statistically significant ($P \leq 0.05$) difference in sensory characteristics between the control yogurt and 0.3% OG yogurt. The control yogurt had the highest acceptability value of 83.26 at 21 d of storage, while 0.3% OG yogurt had the highest acceptability value

of 86.49 at 21 d of storage. Clearly, the addition of 0.3% OG has a positive effect on the acceptability of the yogurt, independently of the storage time.

Table 2. Sensory properties of the control yogurt and 0.3% OG yogurt during storage

Variables	Days	Yogurt samples	
		control yogurt	0.3% OG yogurt
Color	1	8.12 ± 0.24	8.71 ± 0.46*
	7	8.13 ± 0.34*	8.68 ± 0.32**
	14	8.18 ± 0.28*	8.51 ± 0.48**
	21	8.20 ± 0.14*	8.50 ± 0.28*
Structural state	1	26.05 ± 0.47	26.78 ± 0.63**
	7	27.05 ± 0.12*	27.32 ± 0.76*
	14	27.02 ± 0.32*	28.08 ± 0.92**
	21	27.08 ± 0.28*	27.98 ± 0.88*
Texture	1	24.28 ± 0.39	24.97 ± 0.79**
	7	24.92 ± 0.22*	25.12 ± 0.82*
	14	25.04 ± 0.28*	25.98 ± 0.70**
	21	24.88 ± 0.40**	25.88 ± 0.70**
Flavor	1	23.7 ± 0.90	24.67 ± 0.47**
	7	23.1 ± 0.70*	24.02 ± 0.32**
	14	23.0 ± 1.20**	24.12 ± 0.22*
	21	23.1 ± 1.20**	23.88 ± 0.28*

Conclusions

The results of the present work showed that 0.3% oat β -glucan yogurt had good sensory and mouthfeel characteristics by improving viscosity and texture as compared to the control. Due to the lack of dietary fiber in milk and yogurt, β -glucan is a suitable choice for dairy systems.

The 0.3% oat β -glucan yogurt also has high nutritional values as oat β -glucan can act as a prebiotic in probiotic yogurt. Therefore, yogurt containing 0.3% oat β -glucan could be an innovative healthy dairy product for enhancing oat β -glucan consumption.

References

- [1] Chen, M., Ye, X., Shen, D., & Ma, C. (2019). Modulatory effects of gut microbiota on constipation: the commercial beverage yakult shapes stool consistency. *Journal of Neurogastroenterology and Motility*, 25(3), 475-477.
- [2] Xu, S., Gong, Y., Rafique, H., He, T., & Hu, X. (2021). Effect of oat β -glucan addition on the staling properties of wheat-oat blended flour Chinese steamed bread. *Bioactive Carbohydrates and Dietary Fibre*, 26, 100285.
- [3] Sarantis, S.D., Eren, N.M., Kowalczyk, B., Jimenez-Flores, R., & Alvarez, V.B. (2021). Thermodynamic interactions of micellar casein and oat β -glucan in a model food system. *Food Hydrocolloids*, 115, 106559.
- [4] Rosburg, V., Boylston, T., & White, P. (2010). Viability of bifidobacteria strains in yogurt with added oat beta-glucan and corn starch during cold storage. *Journal of Food Science*, 75(5), 439-444.

- [5] Jørgensen, C.E., Abrahamsen, R.K., Rukke, E.O., Hoffmann, T.K., Johansen, A.G., & Skeie, S.B. (2019). Processing of high-protein yoghurt—A review. *International Dairy Journal*, 88, 42-59.
- [6] García-Burgos, M., Moreno-Fernández, J., Alférez, M.J., Díaz-Castro, J., & López-Aliaga, I. (2020). New perspectives in fermented dairy products and their health relevance. *Journal of Functional Foods*, 72, 104059.
- [7] Shiby, V.K., & Mishra, H.N. (2013). Fermented milks and milk products as functional foods—A review. *Critical reviews in food science and nutrition*, 53(5), 482-496.
- [8] Samilyk, M., Helikh, A., Ryzhkova, T., Bolgova, N., & Nazarenko, Y. (2020). Influence of the structure of some types of fillers introduced to the yogurt recipe on changes in its rheological indicators. *Eastern-European Journal of Enterprise Technologies*, 2(11)(104)), 46-51.
- [9] Mudgil, D., Barak, S., & Khatkar, B.S. (2016). Development of functional yoghurt via soluble fiber fortification utilizing enzymatically hydrolyzed guar gum. *Food Bioscience*, 14, 28-33.
- [10] Ibrahim, M., & Selezneva, I.S. (2019). The symbiotic effect of oat β -glucan enriching bio-low fat yogurt. *Scientific journal NRUITMO*, 4, 111-116.
- [11] Ibrahim, M., Barakova, N., & Jōudu, I. (2020). Enrichment of the low-fat yoghurt with oat β -glucan and EPS-producing *Bifidobacterium bifidum* improves its quality. *Agronomy Research*, 18(S3), 1689-1699.
- [12] Qu, X., Nazarenko, Y., Yang, W., Nie, Y., Zhang, Y., & Li, B. (2021). Effect of Oat β -Glucan on the Rheological Characteristics and Microstructure of Set-Type Yogurt. *Molecules*, 26(16), 4752. doi: 10.3390/molecules26164752.
- [13] Aboushanab, S., Vyrova, D., & Selezneva, I. (2018). Characterization of low-and non-fat yogurt manufactured with addition of beta-glucanas a dietary supplement [C]. *AIP Conference Proceedings*. AIP Publishing LLC, 2015(1), 020003.
- [14] Gebara, C., Ribeiro, M. C. E., Chaves, K. S., Gandara, A. L. N., & Gigante, M. L. (2015). Effectiveness of different methodologies for the selective enumeration of *Lactobacillus acidophilus* La5 from yoghurt and Prato cheese. *LWT-Food Science and Technology*, 64(1), 508-513.
- [15] Ladjevardi, Z.S., Yarmand, M.S., Emam-Djomeh, Z., & Niasari-Naslaji, A. (2016). Physicochemical properties and viability of probiotic bacteria of functional synbiotic camel yogurt affected by oat β -glucan during storage. *Journal of agricultural science and technology*, 18(5), 1233-1246.
- [16] Zhao, S., Zhou, Q. Y., Huang, Y.Y., Nan, S.G., & Liu, D.M. (2021). Comparative analysis of physicochemical, rheological, sensory and flavour properties of yoghurts using a new probiotic *Bacillus coagulans* 13002 with traditional yoghurt starter. *International Journal of Food Science & Technology*, 56(4), 1712-1723.
- [17] Deshwal, G.K., Tiwari, S., Kumar, A., Raman, R.K., & Kadyan, S. (2021). Review on factors affecting and control of post-acidification in yoghurt and related products. *Trends in Food Science & Technology*, 109, 499-512.
- [18] Agyemang, P.N., Akonor, P.T., Tortoe, C., Johnsona, P.T., & Manu-Aduening, J. (2020). Effect of the use of starches of three new Ghanaian cassava varieties as a thickener on the physicochemical, rheological and sensory properties of yoghurt. *Scientific African*, 9, e00521.
- [19] Kaur, R., & Riar, C.S. (2020). Sensory, rheological and chemical characteristics during storage of set type full fat yoghurt fortified with barley β -glucan [J]. *Journal of food science and technology*, 57(1), 41-51.

- [20] Westerik, N., Wacoo, A.P., Anyimo, E., Matovu, W., Reid, G., Kort, R., & Sybesma, W. (2019). Improving health and wealth by introduction of an affordable bacterial starter culture for probiotic yoghurt production in Uganda. *Challenges*, 10(1), 2.
- [21] Meybodi, N. M., Mortazavian, A. M., Arab, M., & Nematollahi, A. (2020). Probiotic viability in yoghurt: A review of influential factors. *International Dairy Journal*, 109, 104793.
- [22] Summo, C., De Angelis, D., Difonzo, G., Caponio, F., & Pasqualone, A. (2020). Effectiveness of oat-hull-based ingredient as fat replacer to produce low fat burger with high beta-glucans content. *Foods*, 9(8), 1057.

Вплив зберігання на сенсорні показники, в'язкість та хімічні характеристики йогурту, збагаченого вівсяним β -глюканом

Сяодзинь Цю^{1,2}

¹Сумський національний аграрний університет
4002, вул. Герасима Кондратьєва, 160, м. Суми, Україна

²Хенанський інститут науки і технологій
453003, пр-т. Хуалан, м. Сінсян, Китай

Анотація. Вівсяний β -глюкан має потенціал функціональної добавки до йогурту, однак аналіз сенсорних показників, текстури та хімічні характеристики впродовж зберігання йогурту з додавання цієї добавки потребують більш систематичного вивчення, що зумовлює актуальність дослідження. Метою наукової роботи було проаналізувати доцільність збагачення йогурту 0,3 % вівсяним β -глюканом. Використані методи для отримання нових знань в напрямку розроблення корисних для здоров'я продуктів харчування. З'ясовано, що йогурт із додаванням 0,3 % вівсяного β -глюкану володіє водоутримуючою здатністю. Усі значення рН, під час зберігання, становили від 4,18 до 4,28, що є в межах норми для йогуртів. Встановлено вищу в'язкість експериментального зразка йогурту порівняно з контрольним, а найвищі значення були 58560 ± 2120 ср на 21 добу для йогурту з додаванням 0,3 % вівсяного β -глюкану. Перевірено життєздатність пробіотичних бактерій у йогуртах. За весь період зберігання вміст пробіотиків знижувався, проте йогурт із додаванням 0,3 % вівсяного β -глюкану містив значно більше живих пробіотичних бактерій, порівняно з контролем. Життєздатність пробіотичних бактерій йогурту з додаванням 0,3 % вівсяного β -глюкану на 14 добу ($3,18 \pm 0,2 \times 10^7$ КУО/мл) була лише незначно нижчою, ніж у контрольного йогурту на 1 добу ($3,45 \pm 0,3 \times 10^7$ КУО/мл). Це повністю демонструє, що додавання 0,3 % вівсяного β -глюкану чинить захисну дію на пробіотики в йогурті, що буде корисно для здоров'я людини. Встановлено, що додавання 0,3 % вівсяного β -глюкану, призвело до зниження адгезивності, але посилення твердості та клейкості під час зберігання. Результати сенсорної оцінки показали найвищі значення прийнятності 86,49 на 21 добу зберігання йогурту із додаванням 0,3 % вівсяного β -глюкану. Практична цінність наукової роботи полягає в тому, що розроблений йогурт, який містить 0,3 % вівсяного β -глюкану, може бути інноваційним молочним продуктом для здорового харчування

Ключові слова: йогурт, вівсяний β -глюкан, сенсорні показники, в'язкість, хімічні характеристики